

Bipolar complex fuzzy subalgebras and ideals of BCK/BCI-algebras

Tahir Mahmood^{1,*}, Ubaid ur Rehman¹

¹ Department of Mathematics and Statistics, International Islamic University Islamabad, Pakistan

* Correspondence: tahirbakhat@iiu.edu.pk

Received 09 December 2022

Accepted for publication 3 April 2023

Published 21 April 2023

Abstract

The conception of the bipolar complex fuzzy set (BCFS) is one of the fundamental and significant modifications of the fuzzy set (FS) to tackle the tricky and awkward information. BCFS has a rich and wider structure and has been utilized in various fields. In this article, we introduce the concept of bipolar complex fuzzy (BCF) subalgebras (BCFSAs), BCF ideals (BCFIs) of a BCK/BCI-algebra, along with certain properties. Further, we investigate the relations between BCFSAs and BCFIs and a necessary condition for BCFSAs to be BCFIs. We also investigate characterizations of BCFIs. Moreover, we introduce the notion of equivalence relations on the collection of all BCFIs of BCK/BCI algebra and the associated properties of equivalence relations.

Keywords: bipolar complex fuzzy set, BCI/BCK algebra, bipolar complex fuzzy subalgebra/ideal.

1. Introduction

Every time the non-classical logic gets the benefits and rewards of the conventional logic when tackling the data in numerous parts of vagueness and ambiguity. These days non-conventional logic plays a significant role in computer science because it tackles the data containing fuzziness and vagueness. The conception of BCK/BCI-algebras was investigated by Imai and Iseki (1965) and Iseki (1966) which is considered as the modification of propositional logic. Several people studied the concept of BCK/BCI-algebras and employed it in the environment of FS (Akram and Zhan, 2006; Meng, 2000). The structure of FS was explored by Zadeh in 1965 (Zadeh, 1965) in which the membership grade (MG) of elements span the range of $[0, 1]$. The MG shows how much an element is a part of the FS, if the MG is 1, then it implies that the element belongs entirely to the FS and if the MG is 0, then it implies that the element is not contained in the corresponding FS. Mordeson and Malik (1998) investigated fuzzy commutative algebra, and Dubois and Prade (1979) propounded fuzzy real algebra. The $(\epsilon, \epsilon \vee q)$ and (α, β) fuzzy subalgebras (FSA) in BCI/BCK-algebra was presented by Jun (2005; 2009). FSA of BE-algebras was investigated by Rezaei and Saeid (2011). Akram et al. (2007) explored fuzzy models of $K(G)$ algebras. The fuzzy ideals (FIs) of BCH-algebra were given by Du and Liao (2007). Biswas (1990) investigated fuzzy subgroups and Liu (1982) presented fuzzy invariant subgroups. In the FS, it's hard to indicate the distinction between the contradictory and irrelevant elements and to express the country property of the element. Thus, Zhang (1994) presented the structure of

bipolar FS (BFS) in which the positive MG (PMG) of elements span the range of $[0, 1]$ and negative MG (NMG) of elements span the range of $[-1, 0]$. Akram et al. (2010) explored bipolar fuzzy (BF) K-algebras. Muhiuddin and Al-Kadi (2021) investigated BF implicative ideals in BCK-algebras. Certain characteristics of doubt BF H-ideals in BCI/BCK-algebra were propounded by Al-Masarwah (2018). Muhiuddin (2014) described BF KU-ideals and subalgebras. Muhiuddin et al. (2020) presented novel sort of BF ideals (BFIs) of BCK-algebras. Kawila et al. (2018) explained BF UP-algebras. Lee and Jun (2011) investigated BF a-ideals of BCI-algebras. Jun et al. (2009) explored BF models of certain sorts of ideals in hyper BCK-algebras. Mahmood and Munir (2013) introduced BF subgroups.

There are numerous modifications of the basic conception of FS and complex FS (CFS) is one of them, which was derived by Ramot et al (2022). In CFS, the MG contains both amplitude and phase terms of the elements belonging to $[0, 1]$. Afterward, Tamir et al. (2011) presented another form of CFS. Shagagha (2019) investigated complex fuzzy (CF) lie algebras. Jun and Xin (2019) proposed the application of CFSs in BCK/BCI algebra. Rasuli (2022) explored anti CF lie subalgebras. The other and most advanced modification of FS, BFS, and CFS, is BCFS, investigated by Mahmood and Ur Rehman (2022a) in which the PMG of elements belongs to the first quadrant and NMG of elements belongs to the third quadrant of the unit square. The structure of BCFS is a significant tool for tackling tricky and complicated information and generalizing various prevailing concepts. Due to the wide structure and importance of the BCFS, numerous scholars utilized it in various areas (Mahmood et al., 2021; Rehman et al., 2022; Mahmood and Ur Rehman, 2022b). Yang et al. (2022) introduced the conception of BCF subgroups.

Keeping in view the significance and supremacy of the BCFS, in this study we are going to apply the conception of BCFS to the BCK/BCI-algebra. We investigate the conception of BCF subalgebras and BCF ideals of a BCI/BCK-algebra along with associated properties. Moreover, we develop the relation among BCF subalgebra and BCF ideals and a necessary condition for BCFA to be a BCFI. We also investigate characterizations of BCFI. Further, we propound the conception of equivalence relations on the group of all BCF ideals of BCI/BCK-algebra along with associated properties. The underneath article is managed as, In Section 2, we recalled some fundamental notions related to BCK/BCI-algebra and the notion of BCFS. In Section 3, we investigated the concept of BCFS and BCFI and their related results. The concluding remarks are portrayed in Section 4.

2. Preliminaries

In this Section, we are going to recall some fundamental notions related to BCK/BCI-algebra and the notion of BCFS. The conception of BCI/BCK-algebra is given by Iseki (1966) which has a significant part in the logical algebras and is widely studied by numerous scholars.

Suppose that $\mathfrak{B}(\tau)$ is the collection of all algebras of type $\tau = (2, 0)$. BCI-algebra means the set $(\mathcal{Q}; *, 0) \in \mathfrak{B}(\tau)$ which holds the underneath properties

1. $((q_1 * q_2) * (q_1 * q_3)) * (q_3 * q_2) = 0$ ($\forall q_1, q_2, q_3 \in \mathcal{Q}$)
2. $((q_1 * (q_1 * q_2)) * q_2) = 0$ ($\forall q_1, q_2 \in \mathcal{Q}$)
3. $((q_1 * q_1) = 0)$ ($\forall q_1 \in \mathcal{Q}$)
4. $(q_1 * q_2 = 0, q_2 * q_1 = 0 \Rightarrow q_1 = q_2)$ ($q_1, q_2 \in \mathcal{Q}$)

The BCI-algebra is known as BCK-algebra if it holds the underneath property

5. $(0 * q_1 = 0)$ ($\forall q_1 \in \mathcal{Q}$)

Every BCI/BCK algebra holds the underneath properties

- I. $(q_1 * 0 = 0)$ ($\forall q_1 \in \mathcal{Q}$)
- II. $(q_1 \leq q_2 \Rightarrow q_1 * q_3 \leq q_2 * q_3, q_3 * q_2 \leq q_3 * q_1)$ ($\forall q_1, q_2, q_3 \in \mathcal{Q}$)
- III. $((q_1 * q_2) * q_3 = (q_1 * q_3) * q_2)$ ($\forall q_1, q_2, q_3 \in \mathcal{Q}$)
- IV. $((q_1 * q_3) * (q_2 * q_3) \leq q_1 * q_2)$ ($\forall q_1, q_2, q_3 \in \mathcal{Q}$)

where, $q_1 \leq q_2$ iff $q_1 * q_2 = 0$. Any subset \mathcal{D} of BCI/BCK-algebra \mathcal{Q} is said to be a subalgebra of BCI/BCK-algebra \mathcal{Q} if $q_1 * q_2 \in \mathcal{D} \forall q_1, q_2 \in \mathcal{D}$. A subset \mathfrak{S} is said to be ideal for BCI/BCK-algebra \mathcal{Q} if it contains 0 and if $q_1 * q_2 \in \mathfrak{S}$ and $q_2 \in \mathfrak{S}$, then $q_1 \in \mathfrak{S}$. The ideal \mathfrak{S} of BCI/BCK-algebra \mathcal{Q} holds the underneath axiom

$$q_1 \leq q_2 \text{ and } q_2 \in \mathfrak{S} \Rightarrow q_1 \in \mathfrak{S}$$

An FS θ in BCI/BCK-algebra \mathcal{Q} is considered a fuzzy subalgebra of \mathcal{Q} if $\forall q_1, q_2 \in \mathcal{Q}$

$$\theta(q_1 * q_2) \geq \min(\theta(q_1), \theta(q_2))$$

An FS θ in BCI/BCK-algebra \mathcal{Q} is considered a fuzzy ideal of \mathcal{Q} if $\forall q_1, q_2 \in \mathcal{Q}$

$$\theta(0) \geq \theta(q_1)$$

$$\theta(q_1) \geq \min(\theta(q_1 * q_2), \theta(q_2))$$

The novel structure of BCFS was investigated by Mahmood and Ur Rehman (2022a), given as follow.

The BCFS is of the underneath shape

$$\mathcal{G}_{BCFS} = \left\{ \left(q, \left(\theta_{\mathcal{G}_{BCFS}}^P(q), \theta_{\mathcal{G}_{BCFS}}^N(q) \right) \right) \mid q \in \mathcal{Y} \right\} = \left\{ \left(q, \left(\theta_{\mathcal{G}_{BCFS}}^{RP}(q) + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}(q), \theta_{\mathcal{G}_{BCFS}}^{RN}(q) + \iota \theta_{\mathcal{G}_{BCFS}}^{IN}(q) \right) \right) \mid q \in \mathcal{Q} \right\}$$

where, $\theta_{\mathcal{G}_{BCFS}}^P(q)$ simplifies the positive membership grade and $\theta_{\mathcal{G}_{BCFS}}^N(q)$ simplifies the negative membership grade and $\theta_{\mathcal{G}_{BCFS}}^{RP}(q), \theta_{\mathcal{G}_{BCFS}}^{IP}(q) \in [0, 1], \theta_{\mathcal{G}_{BCFS}}^{RN}(q), \theta_{\mathcal{G}_{BCFS}}^{IN}(q) \in [-1, 0]$. For simplest in this study, we would consider the BCFS as $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$.

3. BCF subalgebras and BCF ideals

Here, we investigate the concept of BCFS and BCFI and their related results. \mathcal{Q} would be considered as BCK/BCI-algebra in this study unless stated otherwise and $0 = (\theta_{0_{BCFS}}^P, \theta_{0_{BCFS}}^N) = (\theta_{0_{BCFS}}^{RP} + \iota \theta_{0_{BCFS}}^{IP}, \theta_{0_{BCFS}}^{RN} + \iota \theta_{0_{BCFS}}^{IN})$ means $0 = (0 + \iota 0, -0 - \iota 0)$ in this article.

Definition 1: A BCFS $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$ in \mathcal{Q} is considered as BCF subalgebra (BCFSA) of \mathcal{Q} if $\forall q_1, q_2 \in \mathcal{Q}$, the underneath holds

1. $\theta_{\mathcal{G}_{BCFS}}^P(q_1 * q_2) \geq \min(\theta_{\mathcal{G}_{BCFS}}^P(q_1), \theta_{\mathcal{G}_{BCFS}}^P(q_2))$ which means that $\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2))$ and $\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2))$
2. $\theta_{\mathcal{G}_{BCFS}}^N(q_1 * q_2) \leq \max(\theta_{\mathcal{G}_{BCFS}}^N(q_1), \theta_{\mathcal{G}_{BCFS}}^N(q_2))$ which means that $\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2))$ and $\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2))$

Example 1: Suppose a BCK-algebra $\mathcal{Q} = \{0, q_1, q_2, q_3\}$ with Cayley table (Table 1) interpreted as

Table 1. Cayley Table of example 1

*	0	q_1	q_2	q_3
0	0	0	0	0
q_1	q_1	0	0	q_1
q_2	q_2	q_1	0	q_2
q_3	q_3	q_3	q_3	0

Now consider a BCFS \mathcal{G}_{BCFS} in \mathcal{Q} such as

$$\mathcal{G}_{BCFS} = \left\{ \left(0, \left(0, 8 + \iota 0.5, -0.6 - \iota 0.4 \right) \right), \left(q_1, \left(0, 8 + \iota 0.5, -0.6 - \iota 0.4 \right) \right), \left(q_2, \left(0, 5 + \iota 0.2, -0.3 - \iota 0.1 \right) \right), \left(q_3, \left(0, 8 + \iota 0.5, -0.6 - \iota 0.4 \right) \right) \right\}$$

Then \mathcal{G}_{BCFS} is a BCFSA of \mathcal{Q} .

Example 2: Suppose a BCI-algebra $\mathcal{Q} = \{0, q, q_1, q_2, q_3\}$ with Cayley table (Table 2) interpreted as

Table 2. Cayley Table of example 2

*	0	q	q ₁	q ₂	q ₃
0	0	0	q ₃	q ₂	q ₁
q	q	0	q ₃	q ₂	q ₁
q ₁	q ₁	q ₁	0	q ₃	q ₂
q ₂	q ₂	q ₂	q ₁	0	q ₃
q ₃	q ₃	q ₃	q ₂	q ₁	0

Now consider a BCFS \mathcal{G}_{BCFS} in \mathcal{Q} such as

$$\mathcal{G}_{BCFS} = \left\{ \left(0, \begin{pmatrix} 0, 9 + \iota 0.6, \\ -0.55 - \iota 0.5 \end{pmatrix} \right), \left(q, \begin{pmatrix} 0, 7 + \iota 0.55, \\ -0.5 - \iota 0.45 \end{pmatrix} \right), \left(q_1, \begin{pmatrix} 0, 4 + \iota 0.3, \\ -0.2 - \iota 0.3 \end{pmatrix} \right), \right. \\ \left. \left(q_2, \begin{pmatrix} 0, 4 + \iota 0.3, \\ -0.2 - \iota 0.3 \end{pmatrix} \right), \left(q_3, \begin{pmatrix} 0, 4 + \iota 0.3, \\ -0.2 - \iota 0.3 \end{pmatrix} \right) \right\}$$

Then \mathcal{G}_{BCFS} is a BCFS of \mathcal{Q} .

Proposition 1: Suppose a BCFS $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$, if \mathcal{G}_{BCFS} is a BCFS of \mathcal{Q} , then $\forall q \in \mathcal{Q} \theta_{\mathcal{G}_{BCFS}}^P(0) \geq \theta_{\mathcal{G}_{BCFS}}^P(q)$ which means that $\theta_{\mathcal{G}_{BCFS}}^{RP}(0) \geq \theta_{\mathcal{G}_{BCFS}}^{RP}(q)$, $\theta_{\mathcal{G}_{BCFS}}^{IP}(0) \geq \theta_{\mathcal{G}_{BCFS}}^{IP}(q)$ and $\theta_{\mathcal{G}_{BCFS}}^N(0) \leq \theta_{\mathcal{G}_{BCFS}}^N(q)$ which means that $\theta_{\mathcal{G}_{BCFS}}^{RN}(0) \leq \theta_{\mathcal{G}_{BCFS}}^{RN}(q)$, $\theta_{\mathcal{G}_{BCFS}}^{IN}(0) \leq \theta_{\mathcal{G}_{BCFS}}^{IN}(q)$.

Proof: Assume that $q \in \mathcal{Q}$, then

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^P(0) = \theta_{\mathcal{G}_{BCFS}}^P(q * q) &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^{RP}(0) = \theta_{\mathcal{G}_{BCFS}}^{RP}(q * q) \text{ and } \theta_{\mathcal{G}_{BCFS}}^{IP}(0) = \theta_{\mathcal{G}_{BCFS}}^{IP}(q * q) \\ &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^{RP}(0) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q), \theta_{\mathcal{G}_{BCFS}}^{RP}(q)) = \theta_{\mathcal{G}_{BCFS}}^{RP}(q) \text{ and} \\ &\theta_{\mathcal{G}_{BCFS}}^{IP}(0) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q), \theta_{\mathcal{G}_{BCFS}}^{IP}(q)) = \theta_{\mathcal{G}_{BCFS}}^{IP}(q) \\ &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^P(0) \geq \theta_{\mathcal{G}_{BCFS}}^P(q) \end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^N(0) = \theta_{\mathcal{G}_{BCFS}}^N(q * q) &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^{RN}(0) = \theta_{\mathcal{G}_{BCFS}}^{RN}(q * q) \text{ and } \theta_{\mathcal{G}_{BCFS}}^{IN}(0) = \theta_{\mathcal{G}_{BCFS}}^{IN}(q * q) \\ &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^{RN}(0) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q), \theta_{\mathcal{G}_{BCFS}}^{RN}(q)) = \theta_{\mathcal{G}_{BCFS}}^{RN}(q) \text{ and} \\ &\theta_{\mathcal{G}_{BCFS}}^{IN}(0) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q), \theta_{\mathcal{G}_{BCFS}}^{IN}(q)) = \theta_{\mathcal{G}_{BCFS}}^{IN}(q) \\ &\Rightarrow \theta_{\mathcal{G}_{BCFS}}^N(0) \leq \theta_{\mathcal{G}_{BCFS}}^N(q). \end{aligned}$$

Definition 2: Consider a BCFS $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$, then

1. The set $\mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) = \{q \in \mathcal{Q} : \theta_{\mathcal{G}_{BCFS}}^{RP}(q) \geq \mathfrak{f} \text{ and } \theta_{\mathcal{G}_{BCFS}}^{IP}(q) \geq \mathfrak{g}\}$ would be positive $(\mathfrak{f}, \mathfrak{g})$ –cut of \mathcal{G}_{BCFS} , where $\mathfrak{f}, \mathfrak{g} \in [0, 1]$.
2. The set $\mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N, (d, e)) = \{q \in \mathcal{Q} : \theta_{\mathcal{G}_{BCFS}}^{RN}(q) \leq d \text{ and } \theta_{\mathcal{G}_{BCFS}}^{IN}(q) \leq e\}$ would be negative (d, e) –cut of \mathcal{G}_{BCFS} , where $d, e \in [-1, 0]$
3. The set $\mathfrak{PN}(\mathcal{G}_{BCFS}, ((\mathfrak{f}, \mathfrak{g}), (d, e))) = \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) \cap \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N, (d, e))$ would be $((\mathfrak{f}, \mathfrak{g}), (d, e))$ –cut of \mathcal{G}_{BCFS} .

Theorem 1: For a BCFS $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ of \mathcal{Q} , the underneath holds

1. $\forall \mathfrak{f}, \mathfrak{g} \in [0, 1] \mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) \neq \emptyset \Rightarrow \mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g}))$ is a subalgebra of \mathcal{Q} .
2. $\forall \mathfrak{d}, \mathfrak{e} \in [-1, 0] \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e})) \neq \emptyset \Rightarrow \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e}))$ is a subalgebra of \mathcal{Q} .

Proof: As $\mathfrak{f}, \mathfrak{g} \in [0, 1]$ and $\mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) \neq \emptyset$. If $q_1, q_2 \in \mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g}))$, $\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \mathfrak{f}$, $\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \mathfrak{g}$, $\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2) \geq \mathfrak{f}$, $\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2) \geq \mathfrak{g}$. Thus $\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \geq \mathfrak{f}$ and $\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \geq \mathfrak{g} \Rightarrow q_1 * q_2 \in \mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g}))$

Hence $\mathfrak{F}(\Theta_{\mathcal{G}_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g}))$ is a subalgebra of \mathcal{Q} .

Now as $\mathfrak{d}, \mathfrak{e} \in [-1, 0]$ and $\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e})) \neq \emptyset$. If $q_1, q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e}))$, $\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \mathfrak{d}$, $\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \mathfrak{e}$, $\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2) \leq \mathfrak{d}$, $\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2) \leq \mathfrak{e}$. Thus $\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1), \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)) \leq \mathfrak{d}$ and $\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)) \leq \mathfrak{e} \Rightarrow q_1 * q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e}))$

Hence $\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (\mathfrak{d}, \mathfrak{e}))$ is a subalgebra of \mathcal{Q} .

Definition 3: A BCFS $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ in \mathcal{Q} is considered as BCF ideal (BCFI) of \mathcal{Q} if $\forall \forall q_1, q_2 \in \mathcal{Q}$, the underneath holds

1. $\Theta_{\mathcal{G}_{BCFS}}^P(0) \geq \Theta_{\mathcal{G}_{BCFS}}^P(q)$ which means that $\Theta_{\mathcal{G}_{BCFS}}^{RP}(0) \geq \Theta_{\mathcal{G}_{BCFS}}^{RP}(q)$ and $\Theta_{\mathcal{G}_{BCFS}}^{IP}(0) \geq \Theta_{\mathcal{G}_{BCFS}}^{IP}(q)$
2. $\Theta_{\mathcal{G}_{BCFS}}^N(0) \leq \Theta_{\mathcal{G}_{BCFS}}^N(q)$ which means that $\Theta_{\mathcal{G}_{BCFS}}^{RN}(0) \leq \Theta_{\mathcal{G}_{BCFS}}^{RN}(q)$ and $\Theta_{\mathcal{G}_{BCFS}}^{IN}(0) \leq \Theta_{\mathcal{G}_{BCFS}}^{IN}(q)$
3. $\Theta_{\mathcal{G}_{BCFS}}^P(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^P(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^P(q_2))$ which means that $\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2))$ and $\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2))$
4. $\Theta_{\mathcal{G}_{BCFS}}^N(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^N(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^N(q_2))$ which means that $\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2))$ and $\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2))$

Example 3: Suppose a BCK-algebra $\mathcal{Q} = \{0, q_1, q_2, q_3, q_4\}$ with Cayley table (Table 3) interpreted as

Table 3. Cayley Table of example 3

*	0	q_1	q_2	q_3	q_4
0	0	0	0	0	0
q_1	q_1	0	q_1	0	0
q_2	q_2	q_2	0	0	0
q_3	q_3	q_3	q_3	0	0
q_4	q_4	q_3	q_4	q_1	0

Now consider a BCFS \mathcal{G}_{BCFS} in \mathcal{Q} such as

$$\mathcal{G}_{BCFS} = \left\{ \left(0, \begin{pmatrix} 0, 67 + \iota 0.5 \\ -0.7 - \iota 0.6 \end{pmatrix} \right), \left(q_1, \begin{pmatrix} 0, 34 + \iota 0.43 \\ -0.4 - \iota 0.5 \end{pmatrix} \right), \left(q_2, \begin{pmatrix} 0, 67 + \iota 0.5 \\ -0.7 - \iota 0.6 \end{pmatrix} \right), \right. \\ \left. \left(q_3, \begin{pmatrix} 0, 34 + \iota 0.43 \\ -0.4 - \iota 0.5 \end{pmatrix} \right), \left(q_4, \begin{pmatrix} 0, 34 + \iota 0.43 \\ -0.4 - \iota 0.5 \end{pmatrix} \right) \right\}$$

Then \mathcal{G}_{BCFS} is a BCFI of \mathcal{Q} .

Proposition 2: Consider a BCFI $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$ of \mathcal{Q} . If in \mathcal{Q} , the inequality $q_1 * q_2 \leq q_3$ holds, then

1. $\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_3))$ and $\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_3))$
2. $\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_2), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_3))$ and $\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q_2), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_3))$

Proof:

1. Assume that $q_1, q_2, q_3 \in \mathcal{Q}$ such that $q_1 * q_2 \leq q_3$, then $(q_1 * q_2) * q_3 = 0$. Thus,

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) &\geq \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \\ &\geq \min(\min(\theta_{\mathcal{G}_{BCFS}}^{RP}((q_1 * q_2) * q_3), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \\ &= \min(\min(\theta_{\mathcal{G}_{BCFS}}^{RP}(0), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \\ &= \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_3)) \end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) &\geq \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \\ &\geq \min(\min(\theta_{\mathcal{G}_{BCFS}}^{IP}((q_1 * q_2) * q_3), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \\ &= \min(\min(\theta_{\mathcal{G}_{BCFS}}^{IP}(0), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \\ &= \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_3)) \end{aligned}$$

2. Assume that $q_1, q_2, q_3 \in \mathcal{Q}$ such that $q_1 * q_2 \leq q_3$, then $(q_1 * q_2) * q_3 = 0$. Thus,

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) &\leq \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)) \\ &\leq \max(\max(\theta_{\mathcal{G}_{BCFS}}^{RN}((q_1 * q_2) * q_3), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)) \\ &= \max(\max(\theta_{\mathcal{G}_{BCFS}}^{RN}(0), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_3)), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)) \\ &= \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_2), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_3)) \end{aligned}$$

and

$$\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2))$$

$$\begin{aligned} &\leq \max\left(\max\left(\theta_{\mathcal{G}_{BCFS}}^{IN}((q_1 * q_2) * q_3), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_3)\right), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)\right) \\ &= \max\left(\max\left(\theta_{\mathcal{G}_{BCFS}}^{IN}(0), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_3)\right), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)\right) \\ &= \max\left(\theta_{\mathcal{G}_{BCFS}}^{IN}(q_2), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_3)\right) \end{aligned}$$

Proposition 3: Consider a BCFI $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$ of \mathcal{Q} . If in \mathcal{Q} , the inequality $q_1 \leq q_2$ holds, then

1. $\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)$ and $\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)$
2. $\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)$ and $\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)$

Proof:

1. Assume that $q_1, q_2, q_3 \in \mathcal{Q}$ such that $q_1 \leq q_2$, then

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) &\geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)\right) \\ &= \min\left(\theta_{\mathcal{G}_{BCFS}}^{RP}(0), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)\right) = \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2) \end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) &\geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)\right) \\ &= \min\left(\theta_{\mathcal{G}_{BCFS}}^{IP}(0), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)\right) = \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2) \end{aligned}$$

2. Assume that $q_1, q_2, q_3 \in \mathcal{Q}$ such that $q_1 \leq q_2$, then

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) &\leq \max\left(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)\right) \\ &= \max\left(\theta_{\mathcal{G}_{BCFS}}^{RN}(0), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)\right) = \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2) \end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) &\leq \max\left(\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)\right) \\ &= \max\left(\theta_{\mathcal{G}_{BCFS}}^{IN}(0), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)\right) = \theta_{\mathcal{G}_{BCFS}}^{IN}(q_2) \end{aligned}$$

Theorem 2: Consider a BCK-algebra \mathcal{Q} , in \mathcal{Q} , each BCFI of \mathcal{Q} is a BCFSA of \mathcal{Q} .

Proof: Assume that $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$ is a BCFI of BCK-algebra of \mathcal{Q} . As $q_1 * q_2 \leq q_1 \forall q_1, q_2 \in \mathcal{Q}$, thus by proposition (3)

$$\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) \geq \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) \geq \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \text{ and}$$

$$\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2) \leq \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1), \theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2) \leq \theta_{\mathcal{G}_{BCFS}}^{IN}(q_1)$$

and from Def (3), we have

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) &\geq \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)\right) \\ &\geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1), \theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)\right) \end{aligned}$$

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) &\geq \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)\right) \\ &\geq \min\left(\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1), \theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)\right) \end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2) &\leq \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \max\left(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)\right) \\ &\leq \max\left(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)\right) \end{aligned}$$

$$\begin{aligned}\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2) &\leq \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)) \\ &\leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2))\end{aligned}$$

Therefore, \mathcal{G}_{BCFS} is a BCFS of \mathcal{Q} .

Remark 1: The converse of the above Theorem usually does not hold.

Example 4: Consider example 1 in which $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ is a BCFS of \mathcal{Q} but \mathcal{G}_{BCFS} is not a BCFI of \mathcal{Q} because

$$\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2) = -0.3 > -0.6 = \max(\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2 * q_1), \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1))$$

The following result would show with what condition a BCFS can be a BCFI in BCK-algebra.

Theorem 3: Suppose a BCFS $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ of BCK-algebra \mathcal{Q} such that $\forall q_1, q_2, q_3 \in \mathcal{Q}$ $\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_3))$ and $\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_3))$, $\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2), \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_3))$ and $\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_3))$ holds and satisfies the inequality $q_1 * q_2 \leq q_3$, then \mathcal{G}_{BCFS} is a BCFI of \mathcal{Q} .

Proof: From Proposition (1) we have that $\forall q \in \mathcal{Q}$

$$\Theta_{\mathcal{G}_{BCFS}}^{RP}(0) \geq \Theta_{\mathcal{G}_{BCFS}}^{RP}(q), \Theta_{\mathcal{G}_{BCFS}}^{IP}(0) \geq \Theta_{\mathcal{G}_{BCFS}}^{IP}(q) \text{ and}$$

$$\Theta_{\mathcal{G}_{BCFS}}^{RN}(0) \leq \Theta_{\mathcal{G}_{BCFS}}^{RN}(q), \Theta_{\mathcal{G}_{BCFS}}^{IN}(0) \leq \Theta_{\mathcal{G}_{BCFS}}^{IN}(q)$$

As $q_1 * (q_1 * q_2) \leq q_2 \forall q_1, q_2 \in \mathcal{Q}$, so by Proposition (2)

$$\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2))$$

and

$$\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \leq \max(\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2))$$

This implies that \mathcal{G}_{BCFS} is a BCFI of \mathcal{Q} .

Theorem 4: Consider a BCFS $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ in \mathcal{Q} , then \mathcal{G}_{BCFS} is said to be BCFI of \mathcal{Q} iff the underneath holds

- $\forall f, g \in [0, 1] \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g)) \neq \emptyset \Rightarrow \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g))$ is an ideal of \mathcal{Q} .
- $\forall d, e \in [-1, 0] \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e)) \neq \emptyset \Rightarrow \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e))$ is an ideal of \mathcal{Q} .

Proof: Let $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ be a BCFI of \mathcal{Q} and $f, g \in [0, 1]$, $d, e \in [-1, 0]$ such that $\mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g)) \neq \emptyset$ and $\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e)) \neq \emptyset$. $0 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g)) \cap \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e))$. Suppose that $q_1, q_2 \in \mathcal{Q}$ such that $q_1 * q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g))$ and $q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P, (f, g))$ and suppose that $q_3, q_4 \in \mathcal{Q}$ such that $q_3 * q_4 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e))$ and $q_4 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e))$, then

$$\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) \geq f, \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2) \geq f, \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) \geq g, \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2) \geq g \text{ and}$$

$$\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_3 * q_4) \leq d, \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_4) \leq d, \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_3 * q_4) \leq e, \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_4) \leq e$$

By Def (3)

$$\begin{aligned}\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \geq f, \\ \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \geq g\end{aligned}$$

and

$$\begin{aligned} \theta_{\mathcal{G}_{BCFS}}^{RN}(q_3) &\leq \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q_3 * q_4), \theta_{\mathcal{G}_{BCFS}}^{RN}(q_4)) \leq d, \\ \theta_{\mathcal{G}_{BCFS}}^{RI}(q_3) &\leq \max(\theta_{\mathcal{G}_{BCFS}}^{RI}(q_3 * q_4), \theta_{\mathcal{G}_{BCFS}}^{RI}(q_4)) \leq e \end{aligned}$$

$\Rightarrow q_1 \in \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}))$ and $q_3 \in \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e))$. Thus, $\mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}))$ and $\mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e))$ are ideals of \mathcal{Q} .

Conversely, assume that conditions 1 and 2 are valid. Suppose that $\theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) = \mathfrak{f}$, $\theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) = \mathfrak{g}$, $\theta_{\mathcal{G}_{BCFS}}^{RN}(q_1) = d$, and $\theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) = e$, for any $q_1 \in \mathcal{Q}$, then $q_1 \in \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g})) \cap \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e)) \Rightarrow \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g})) \neq \emptyset$ and $\mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e)) \neq \emptyset$. As $\mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}))$ and $\mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e))$ are ideals of \mathcal{Q} , $0 \in \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g})) \cap \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d, e))$. Thus, $\theta_{\mathcal{G}_{BCFS}}^{RP}(0) \geq \mathfrak{f} = \theta_{\mathcal{G}_{BCFS}}^{RP}(q_1)$, $\theta_{\mathcal{G}_{BCFS}}^{IP}(0) \geq \mathfrak{g} = \theta_{\mathcal{G}_{BCFS}}^{IP}(q_1)$, $\theta_{\mathcal{G}_{BCFS}}^{RN}(0) \leq d = \theta_{\mathcal{G}_{BCFS}}^{RN}(q_1)$ and $\theta_{\mathcal{G}_{BCFS}}^{IN}(0) \leq e = \theta_{\mathcal{G}_{BCFS}}^{IN}(q_1) \forall q_1 \in \mathcal{Q}$. If $\exists q'_1, q'_2, q'_3, q'_4 \in \mathcal{Q}$ such that

$$\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1) < \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q'_2)), \theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1) < \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q'_2))$$

and

$$\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3) > \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{RN}(q'_4)), \theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3) > \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{IN}(q'_4))$$

then by taking

$$\mathfrak{f}_0 = \frac{1}{2}(\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1) + \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q'_2)))$$

$$\mathfrak{g}_0 = \frac{1}{2}(\theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1) + \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q'_2)))$$

and

$$d_0 = \frac{1}{2}(\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3) + \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{RN}(q'_4)))$$

$$e_0 = \frac{1}{2}(\theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3) + \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{IN}(q'_4)))$$

We have

$$\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1) < \mathfrak{f}_0 < \min(\theta_{\mathcal{G}_{BCFS}}^{RP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{RP}(q'_2))$$

$$\theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1) < \mathfrak{g}_0 < \min(\theta_{\mathcal{G}_{BCFS}}^{IP}(q'_1 * q'_2), \theta_{\mathcal{G}_{BCFS}}^{IP}(q'_2))$$

and

$$\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3) > d_0 > \max(\theta_{\mathcal{G}_{BCFS}}^{RN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{RN}(q'_4))$$

$$\theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3) > e_0 > \max(\theta_{\mathcal{G}_{BCFS}}^{IN}(q'_3 * q'_4), \theta_{\mathcal{G}_{BCFS}}^{IN}(q'_4))$$

Thus, $q'_1 \notin \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}_0, \mathfrak{g}_0))$, $q'_1 * q'_2 \in \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}_0, \mathfrak{g}_0))$, $q'_2 \in \mathfrak{P}(\theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}_0, \mathfrak{g}_0))$, $q'_3 \notin \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d_0, e_0))$, $q'_3 * q'_4 \in \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d_0, e_0))$, $q'_4 \in \mathfrak{N}(\theta_{\mathcal{G}_{BCFS}}^N(d_0, e_0))$ which is a contradiction and therefore, \mathcal{G}_{BCFS} is BCFI of \mathcal{Q} .

Corollary 1: For a BCFI $\mathcal{G}_{BCFS} = (\theta_{\mathcal{G}_{BCFS}}^P, \theta_{\mathcal{G}_{BCFS}}^N) = (\theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \theta_{\mathcal{G}_{BCFS}}^{IP}, \theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \theta_{\mathcal{G}_{BCFS}}^{IN})$ of \mathcal{Q} , the intersection of a non-empty $(\mathfrak{f}, \mathfrak{g})$ –cut and a non-empty (d, e) –cut of \mathcal{G}_{BCFS} is an ideal of \mathcal{Q} .

Remarks 2:

1. The union of a non-empty $(\mathfrak{f}, \mathfrak{g})$ –cut and a non-empty (d, e) –cut of \mathcal{G}_{BCFS} may not be an ideal of \mathcal{Q}

Example 5: Suppose a BCI-algebra $\mathcal{Q} = \{0, q_1, q_2, q_3\}$ with Cayley table (Table 4) interpreted as

Table 4. Cayley Table of example 5

*	0	q ₁	q ₂	q ₃
0	0	q ₁	q ₂	q ₃
q ₁	q ₁	0	q ₃	q ₂
q ₂	q ₂	q ₃	0	q ₁
q ₃	q ₃	q ₂	q ₁	0

Suppose a BCFS in \mathcal{Q} as

$$\mathcal{G}_{BCFS} = \left\{ \left(0, \begin{pmatrix} 0, 6 + \iota 0.4 \\ -0.8 - \iota 0.5 \end{pmatrix} \right), \left(q_1, \begin{pmatrix} 0, 5 + \iota 0.3 \\ -0.4 - \iota 0.3 \end{pmatrix} \right), \left(q_2, \begin{pmatrix} 0, 4 + \iota 0.2 \\ -0.6 - \iota 0.4 \end{pmatrix} \right), \left(q_3, \begin{pmatrix} 0, 4 + \iota 0.2 \\ -0.3 - \iota 0.1 \end{pmatrix} \right) \right\}$$

Then,

$$\mathfrak{P} \left(\Theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}) \right) = \begin{cases} \emptyset & \text{if } 0.6 < \mathfrak{f} \leq 1, \text{ and } 0.4 < \mathfrak{g} \leq 1 \\ \{0\} & \text{if } 0.5 < \mathfrak{f} \leq 0.6, \text{ and } 0.3 < \mathfrak{g} \leq 0.4 \\ \{0, q_1\} & \text{if } 0.4 < \mathfrak{f} \leq 0.5, \text{ and } 0.2 < \mathfrak{g} \leq 0.3 \\ \mathcal{Q} & \text{if } 0 \leq \mathfrak{f} \leq 0.4, \text{ and } 0 \leq \mathfrak{g} \leq 0.2, \end{cases}$$

and

$$\mathfrak{N} \left(\Theta_{\mathcal{G}_{BCFS}}^N(d, e) \right) = \begin{cases} \emptyset & \text{if } -1 \leq d < -0.8, \text{ and } -1 \leq e < -0.5 \\ \{0\} & \text{if } -0.8 \leq d < -0.6, \text{ and } -0.5 \leq e < -0.4 \\ \{0, q_2\} & \text{if } -0.6 \leq d < -0.4, \text{ and } -0.4 \leq e < -0.1 \\ \mathcal{Q} & \text{if } -0.4 \leq d < 0, \text{ and } -0.1 \leq e < 0, \end{cases}$$

It is clear from Theorem (4) that \mathcal{G}_{BCFS} is a BCFI of \mathcal{Q} . But $\mathfrak{P} \left(\Theta_{\mathcal{G}_{BCFS}}^P(0.5, 0.4) \right) \cup \mathfrak{N} \left(\Theta_{\mathcal{G}_{BCFS}}^N(-0.6, -0.4) \right) = \{0, q_1\} \cup \{0, q_2\} = \{0, q_1, q_2\}$ and not a BCFI of \mathcal{Q} because $q_3 * q_1 = q_2 \in \{0, q_1, q_2\}$ but $q_3 \notin \{0, q_1, q_2\}$.

2. If $d = -\mathfrak{f}$ and $e = -\mathfrak{g}$, then also the union of a non-empty $\mathfrak{P} \left(\Theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}) \right)$ and a non-empty $\mathfrak{N} \left(\Theta_{\mathcal{G}_{BCFS}}^N(-\mathfrak{f}, -\mathfrak{g}) \right)$ of \mathcal{G}_{BCFS} is not an ideal of \mathcal{Q} .

Example 6 Suppose a BCI-algebra $\mathcal{Q} = \{0, 1, q_1, q_2, q_3\}$ with Cayley table (Table 5) interpreted as

Table 5. Cayley Table of example 6

*	0	1	q ₁	q ₂	q ₃
0	0	0	q ₁	q ₂	q ₃
1	1	0	q ₁	q ₂	q ₃
q ₁	q ₁	q ₁	0	q ₃	q ₂
q ₂	q ₂	q ₂	q ₃	0	q ₁
q ₃	q ₃	q ₃	q ₂	q ₁	0

Suppose a BCFS in \mathcal{Q} as

$$\mathcal{G}_{BCFS} = \left\{ \left(0, \begin{pmatrix} 0, 9 + \iota 0.6 \\ -0.6 - \iota 0.5 \end{pmatrix} \right), \left(1, \begin{pmatrix} 0, 7 + \iota 0.4 \\ -0.6 - \iota 0.5 \end{pmatrix} \right), \left(q_1, \begin{pmatrix} 0, 4 + \iota 0.3 \\ -0.3 - \iota 0.2 \end{pmatrix} \right), \right. \\ \left. \left(q_2, \begin{pmatrix} 0, 2 + \iota 0.1 \\ -0.4 - \iota 0.3 \end{pmatrix} \right), \left(q_3, \begin{pmatrix} 0, 2 + \iota 0.1 \\ -0.3 - \iota 0.2 \end{pmatrix} \right) \right\}$$

Then,

$$\mathfrak{P} \left(\Theta_{\mathcal{G}_{BCFS}}^P(\mathfrak{f}, \mathfrak{g}) \right) = \begin{cases} \emptyset & \text{if } 0.9 < \mathfrak{f} \leq 1, \text{ and } 0.6 < \mathfrak{g} \leq 1 \\ \{0\} & \text{if } 0.7 < \mathfrak{f} \leq 0.9, \text{ and } 0.4 < \mathfrak{g} \leq 0.6 \\ \{0, 1\} & \text{if } 0.4 < \mathfrak{f} \leq 0.7, \text{ and } 0.3 < \mathfrak{g} \leq 0.4 \\ \{0, 1, q_1\} & \text{if } 0.2 < \mathfrak{f} \leq 0.4, \text{ and } 0.1 < \mathfrak{g} \leq 0.3 \\ \mathcal{Q} & \text{if } 0 \leq \mathfrak{f} \leq 0.2, \text{ and } 0 \leq \mathfrak{g} \leq 0.1, \end{cases}$$

and

$$\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(d, e)) = \begin{cases} \emptyset & \text{if } -1 \leq d < -0.6, \text{ and } -1 \leq e < -0.5 \\ \{0, 1\} & \text{if } -0.6 \leq d < -0.4, \text{ and } -0.5 \leq e < -0.3 \\ \{0, 1, q_2\} & \text{if } -0.4 \leq d < -0.3, \text{ and } -0.3 \leq e < -0.2 \\ \mathcal{Q} & \text{if } -0.3 \leq d < 0, \text{ and } -0.2 \leq e < 0, \end{cases}$$

It is clear from Theorem (4) that \mathcal{G}_{BCFS} is a BCFI of \mathcal{Q} . But $\mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(0.4, 0.3)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-0.4, -0.3)) = \{0, 1, q_1\} \cup \{0, 1, q_2\} = \{0, 1, q_1, q_2\}$ and not a BCFI of \mathcal{Q} because $q_3 * q_1 = q_2 \in \{0, 1, q_1, q_2\}$ but $q_3 \notin \{0, q_1, q_2\}$.

Theorem 5: If $\mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN})$ is a BCFI of \mathcal{Q} such that $\forall q \in \mathcal{Q}$,

$$\Theta_{\mathcal{G}_{BCFS}}^{RP}(q) + \Theta_{\mathcal{G}_{BCFS}}^{RN}(q) \geq 0 \quad \text{and} \quad \Theta_{\mathcal{G}_{BCFS}}^{IP}(q) + \Theta_{\mathcal{G}_{BCFS}}^{IN}(q) \geq 0 \quad (1)$$

then, the union of a non-empty (f, g) -cut and a non-empty $(-f, -g)$ -cut (i.e. $d = -f, e = -g$) of \mathcal{G}_{BCFS} is an ideal of \mathcal{Q} .

Proof: Suppose $f, g \in [0, 1]$ and as $\mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \neq \emptyset$ and $\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g)) \neq \emptyset$, are the ideal of \mathcal{Q} by employing Theorem (4). Thus, $0 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cap \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$. Suppose that $q_1, q_2 \in \mathcal{Q}$ such that $q_1 * q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$ and $q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$. From this, we get the four cases

- i. $q_1 * q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g))$ and $q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g))$
- ii. $q_1 * q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g))$ and $q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$
- iii. $q_1 * q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$ and $q_2 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g))$
- iv. $q_1 * q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$ and $q_2 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$

(i) implies that $q_1 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \subseteq \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$, (iv) implies that $q_1 \in \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g)) \subseteq \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g))$. For (ii), we have

$$\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2) \geq f, \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2) \geq g \quad \text{and} \quad \Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2) \leq -f, \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2) \leq -g$$

So from Def (3) and Eq. (1), we have

$$\begin{aligned} \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), -\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_2)) \geq f \\ \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), -\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_2)) \geq g \\ &\Rightarrow q_1 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \subseteq \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g)) \end{aligned}$$

now (iii) implies that

$$\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2) \leq -f, \Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2) \leq -g \quad \text{and} \quad \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2) \geq f, \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2) \geq g$$

So from Def (3) and Eq. (1), we have

$$\begin{aligned} \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{RP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \geq \min(-\Theta_{\mathcal{G}_{BCFS}}^{RN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{RP}(q_2)) \geq f \\ \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1) &\geq \min(\Theta_{\mathcal{G}_{BCFS}}^{IP}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \geq \min(-\Theta_{\mathcal{G}_{BCFS}}^{IN}(q_1 * q_2), \Theta_{\mathcal{G}_{BCFS}}^{IP}(q_2)) \geq g \\ &\Rightarrow q_1 \in \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \subseteq \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g)). \quad \text{Thus, } \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS}}^P(f, g)) \cup \\ &\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N(-f, -g)) \text{ is an ideal of } \mathcal{Q}. \end{aligned}$$

Suppose that $BCFI(\mathcal{Q})$ is the group of all BCFIs of \mathcal{Q} , $f, g \in [0, 1]$ and $d, e \in [-1, 0]$, then we introduce binary relations $\mathfrak{RP}^{f, g}$ and $\mathfrak{RN}^{d, e}$ on $BCFI(\mathcal{Q})$ as

$$(\mathcal{G}_{BCFS-1}, \mathcal{G}_{BCFS-2}) \in \mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}} \Leftrightarrow \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS-1}}^P, (\mathfrak{f}, \mathfrak{g})) = \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS-2}}^P, (\mathfrak{f}, \mathfrak{g}))$$

$$(\mathcal{G}_{BCFS-1}, \mathcal{G}_{BCFS-2}) \in \mathfrak{RN}^{d, e} \Leftrightarrow \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS-1}}^N, (d, e)) = \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS-2}}^N, (d, e))$$

respectively, $\forall \mathcal{G}_{BCFS-1} = (\Theta_{\mathcal{G}_{BCFS-1}}^P, \Theta_{\mathcal{G}_{BCFS-1}}^N) = (\Theta_{\mathcal{G}_{BCFS-1}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IP}, \Theta_{\mathcal{G}_{BCFS-1}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IN})$ and $\mathcal{G}_{BCFS-2} = (\Theta_{\mathcal{G}_{BCFS-2}}^P, \Theta_{\mathcal{G}_{BCFS-2}}^N) = (\Theta_{\mathcal{G}_{BCFS-2}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS-2}}^{IP}, \Theta_{\mathcal{G}_{BCFS-2}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS-2}}^{IN}) \in BCFI(Q)$. Obviously, $\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ and $\mathfrak{RN}^{d, e}$ are equivalence relations on $BCFI(Q)$. For $\mathcal{G}_{BCFS-1} = (\Theta_{\mathcal{G}_{BCFS-1}}^P, \Theta_{\mathcal{G}_{BCFS-1}}^N) = (\Theta_{\mathcal{G}_{BCFS-1}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IP}, \Theta_{\mathcal{G}_{BCFS-1}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IN}) \in BCFI(Q)$, suppose that $[\mathcal{G}_{BCFS-1}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}}$ ($[\mathcal{G}_{BCFS-1}]_{\mathfrak{RN}^{d, e}}$) is an equivalence class of \mathcal{G}_{BCFS-1} modular $\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ ($\mathfrak{RN}^{d, e}$). Further, we would signify the collection of all equivalence classes modular $\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ ($\mathfrak{RN}^{d, e}$) by $BCFI(Q)/\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ ($BCFI(Q)/\mathfrak{RN}^{d, e}$) i.e.

$$BCFI(Q)/\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}} = \{[\mathcal{G}_{BCFS-1}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}} \mid \mathcal{G}_{BCFS-1} \in BCFI(Q)\}$$

$$BCFI(Q)/\mathfrak{RN}^{d, e} = \{[\mathcal{G}_{BCFS-1}]_{\mathfrak{RN}^{d, e}} \mid \mathcal{G}_{BCFS-1} \in BCFI(Q)\}$$

Next, Suppose that the collection of all ideals of Q is signified by $Ideal(Q)$, define maps

$$\Gamma_{(\mathfrak{f}, \mathfrak{g})}: BCFI(Q) \rightarrow Ideal(Q) \cup \{\emptyset\} \text{ by } \Gamma_{(\mathfrak{f}, \mathfrak{g})}(\mathcal{G}_{BCFS}) = \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS-1}}^P, (\mathfrak{f}, \mathfrak{g}))$$

and

$$\Gamma_{(d, e)}: BCFI(Q) \rightarrow Ideal(Q) \cup \{\emptyset\} \text{ by } \Gamma_{(d, e)}(\mathcal{G}_{BCFS}) = \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS-2}}^N, (d, e))$$

$\forall \mathcal{G}_{BCFS} = (\Theta_{\mathcal{G}_{BCFS}}^P, \Theta_{\mathcal{G}_{BCFS}}^N) = (\Theta_{\mathcal{G}_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IP}, \Theta_{\mathcal{G}_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS}}^{IN}) \in BCFI(Q)$. Clearly $\Gamma_{(\mathfrak{f}, \mathfrak{g})}$ and $\Gamma_{(d, e)}$ are well defined.

Theorem 6: Let $\mathfrak{f}, \mathfrak{g} \in (0, 1]$ and $d, e \in [-1, 0)$, then maps $\Gamma_{(\mathfrak{f}, \mathfrak{g})}$ and $\Gamma_{(d, e)}$ are surjective.

Proof: Clearly a BCFS $0 = (\Theta_{0_{BCFS}}^P, \Theta_{0_{BCFS}}^N) = (\Theta_{0_{BCFS}}^{RP} + \iota \Theta_{0_{BCFS}}^{IP}, \Theta_{0_{BCFS}}^{RN} + \iota \Theta_{0_{BCFS}}^{IN})$ is a BCFI of Q , where $\forall q \in 0, \Theta_{0_{BCFS}}^{RP}(q) = \Theta_{0_{BCFS}}^{IP}(q) = \Theta_{0_{BCFS}}^{RN}(q) = \Theta_{0_{BCFS}}^{IN}(q) = 0$. Then

$$\Gamma_{(\mathfrak{f}, \mathfrak{g})}(0) = \mathfrak{P}(\Theta_{0_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) = \emptyset$$

and

$$\Gamma_{(d, e)}(0) = \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS}}^N, (d, e)) = \emptyset$$

now for any non-empty $\mathcal{H} \in Ideal(Q)$, suppose a BCFS $\mathcal{G}(\mathcal{H})_{BCFS} = (\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P, \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N) = (\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IP}, \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IN})$ in Q , and

$$\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P: Q \rightarrow [0, 1] + \iota [0, 1] \text{ defined as } \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P = \begin{cases} 1 + \iota 1 & \text{if } q \in \mathcal{H} \\ 0 + \iota 0 & \text{otherwise} \end{cases}$$

and

$$\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N: Q \rightarrow [-1, 0] + \iota [-1, 0] \text{ defined as } \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N = \begin{cases} -1 - \iota 1 & \text{if } q \in \mathcal{H} \\ -0 - \iota 0 & \text{otherwise} \end{cases}$$

Then clearly $\mathcal{G}(\mathcal{H})$ is a BCFI of Q . Next, we have

$$\Gamma_{(\mathfrak{f}, \mathfrak{g})}(\mathcal{G}(\mathcal{H})) = \mathfrak{P}(\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) = \{q \in Q \mid \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RP}(q) \geq \mathfrak{f} \ \& \ \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IP}(q) \geq \mathfrak{g}\}$$

$$= \{q \in Q \mid \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RP}(q) = 1 \ \& \ \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IP}(q) = 1\} = \mathcal{H}$$

and

$$\Gamma_{(d, e)}(\mathcal{G}(\mathcal{H})) = \mathfrak{N}(\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N, (d, e)) = \{q \in Q \mid \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RN}(q) \leq d \ \& \ \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IN}(q) \leq e\}$$

$$\{q \in Q \mid \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RN}(q) = -1 \ \& \ \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IN}(q) = -1\} = \mathcal{H}$$

Thus, $\Gamma_{(\mathfrak{f}, \mathfrak{g})}$ and $\Gamma_{(d, e)}$ are onto (surjective).

Theorem 7: For every $\mathfrak{f}, \mathfrak{g} \in (0, 1]$ and $d, e \in [-1, 0)$, the quotient sets $BCFI(Q)/\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ and $BCFI(Q)/\mathfrak{RN}^{d, e}$ are equipotent to $Ideal(Q) \cap \{\emptyset\}$.

Proof: Define two maps

$$\Gamma'_{(\mathfrak{f}, \mathfrak{g})}: BCFI(Q)/\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}} \rightarrow Ideal(Q) \cup \{\emptyset\} \text{ as } \Gamma'_{(\mathfrak{f}, \mathfrak{g})}([\mathcal{G}_{BCFS}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}}) = \Gamma_{(\mathfrak{f}, \mathfrak{g})}(\mathcal{G}_{BCFS})$$

and

$$\Gamma'_{(d, e)}: BCFI(Q)/\mathfrak{RN}^{d, e} \rightarrow Ideal(Q) \cup \{\emptyset\} \text{ as } \Gamma'_{(d, e)}([\mathcal{G}_{BCFS}]_{\mathfrak{RN}^{d, e}}) = \Gamma_{(d, e)}(\mathcal{G}_{BCFS})$$

For $\mathfrak{f}, \mathfrak{g} \in (0, 1]$ and $d, e \in [-1, 0)$ and $\forall \mathcal{G}_{BCFS} \in BCFI(Q)$. Now each $\mathcal{G}_{BCFS-1} = (\Theta_{\mathcal{G}_{BCFS-1}}^P, \Theta_{\mathcal{G}_{BCFS-1}}^N) = (\Theta_{\mathcal{G}_{BCFS-1}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IP}, \Theta_{\mathcal{G}_{BCFS-1}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS-1}}^{IN})$ and $\mathcal{G}_{BCFS-2} = (\Theta_{\mathcal{G}_{BCFS-2}}^P, \Theta_{\mathcal{G}_{BCFS-2}}^N) = (\Theta_{\mathcal{G}_{BCFS-2}}^{RP} + \iota \Theta_{\mathcal{G}_{BCFS-2}}^{IP}, \Theta_{\mathcal{G}_{BCFS-2}}^{RN} + \iota \Theta_{\mathcal{G}_{BCFS-2}}^{IN}) \in BCFI(Q)$, if $\mathfrak{P}(\Theta_{\mathcal{G}_{BCFS-1}}^P, (\mathfrak{f}, \mathfrak{g})) = \mathfrak{P}(\Theta_{\mathcal{G}_{BCFS-2}}^P, (\mathfrak{f}, \mathfrak{g}))$ and $\mathfrak{N}(\Theta_{\mathcal{G}_{BCFS-1}}^N, (d, e)) = \mathfrak{N}(\Theta_{\mathcal{G}_{BCFS-2}}^N, (d, e))$, then $(\mathcal{G}_{BCFS-1}, \mathcal{G}_{BCFS-2}) \in \mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}$ and $(\mathcal{G}_{BCFS-1}, \mathcal{G}_{BCFS-2}) \in \mathfrak{RN}^{d, e}$, thus $[\mathcal{G}_{BCFS-1}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}} = [\mathcal{G}_{BCFS-2}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}}$ and $[\mathcal{G}_{BCFS-1}]_{\mathfrak{RN}^{d, e}} = [\mathcal{G}_{BCFS-2}]_{\mathfrak{RN}^{d, e}}$. Consequently, $\Gamma'_{(\mathfrak{f}, \mathfrak{g})}$ and $\Gamma'_{(d, e)}$ are one-one (injective). For non-empty $\mathcal{H} \in Ideal(Q)$, suppose a BCFS $\mathcal{G}(\mathcal{H})_{BCFS} = (\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P, \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N) = (\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RP} + \iota \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IP}, \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{RN} + \iota \Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^{IN})$ in Q presented in Theorem (6), then

$$\Gamma'_{(\mathfrak{f}, \mathfrak{g})}([\mathcal{G}(\mathcal{H})_{BCFS}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}}) = \Gamma_{(\mathfrak{f}, \mathfrak{g})}(\mathcal{G}(\mathcal{H})_{BCFS}) = \mathfrak{P}(\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) = \mathcal{H}$$

and

$$\Gamma'_{(d, e)}([\mathcal{G}(\mathcal{H})_{BCFS}]_{\mathfrak{RN}^{d, e}}) = \Gamma_{(d, e)}(\mathcal{G}(\mathcal{H})_{BCFS}) = \mathfrak{N}(\Theta_{\mathcal{G}(\mathcal{H})_{BCFS}}^N, (d, e)) = \mathcal{H}$$

Next, For BCFI $0 = (\Theta_{0_{BCFS}}^P, \Theta_{0_{BCFS}}^N) = (\Theta_{0_{BCFS}}^{RP} + \iota \Theta_{0_{BCFS}}^{IP}, \Theta_{0_{BCFS}}^{RN} + \iota \Theta_{0_{BCFS}}^{IN})$, we have

$$\Gamma'_{(\mathfrak{f}, \mathfrak{g})}([0_{BCFS}]_{\mathfrak{RP}^{\mathfrak{f}, \mathfrak{g}}}) = \Gamma_{(\mathfrak{f}, \mathfrak{g})}(0_{BCFS}) = \mathfrak{P}(\Theta_{0_{BCFS}}^P, (\mathfrak{f}, \mathfrak{g})) = \emptyset$$

and

$$\Gamma'_{(d, e)}([0_{BCFS}]_{\mathfrak{RN}^{d, e}}) = \Gamma_{(d, e)}(0_{BCFS}) = \mathfrak{N}(\Theta_{0_{BCFS}}^N, (d, e)) = \emptyset$$

Hence, $\Gamma'_{(\mathfrak{f}, \mathfrak{g})}$ and $\Gamma'_{(d, e)}$ are surjective (onto) and the required proof is completed.

4. Conclusion

We developed this study by keeping in view the importance and significance of the BCFS theory as BCFS theory is one of the richest and most modified theories in the prevailing literature. This article contained the conception of BCFSAs and BCFIs of BCK/BCI-algebras with various properties. Also, this article contained the relations between BCFSAs and a BCFI and a significant condition for BCFSAs to be a BCFI. Furthermore, we proposed characterizations of BCFI in this study. At last, this study contained the conception of equivalence relations on the group of all BCFIs of BCK/BCI-algebra and the linked properties.

In the future, we would like to employ the investigated work to some of the prevailing works like BCF soft sets (Mahmood et al, 2022), bipolar complex intuitionistic FSs (Al-Husban, 2022; Jan, 2022).

References:

Akram, M., & Zhan, J. (2006). On sensible fuzzy ideals of BCK-algebras with respect to a t-conorm. International journal of mathematics and mathematical sciences, 2006.

Akram, M., Dar, K. H., Jun, Y. B., & Roh, E. H. (2007). Fuzzy structures of K (G)-algebra. Southeast Asian Bulletin of Mathematics, 31(4), 625-637.

Akram, M., Saeid, A. B., Shum, K. P., & Meng, B. L. (2010). Bipolar fuzzy K-algebras. International Journal of Fuzzy Systems, 12(3), 252-258.

Al-Husban, A. (2022). Bipolar Complex Intuitionistic Fuzzy Sets. Earthline Journal of Mathematical Sciences, 8(2), 273-280.

Al-Masarwah, A. M. (2018). On some properties of doubt bipolar fuzzy H-ideals in BCK/BCI-algebras. European journal of pure and applied mathematics, 11(3), 652-670.

Biswas, R. (1990). Fuzzy subgroups and anti fuzzy subgroups. Fuzzy sets and systems, 35(1), 121-124.

- Du, J., & Liao, Z. (2007, August). _Generalized Fuzzy Ideals of BCH-Algebra. In Fourth International Conference on Fuzzy Systems and Knowledge Discovery (FSKD 2007), IEEE, 1, 294-299.
- Dubois, D., & Prade, H. (1979). Fuzzy real algebra: some results. *Fuzzy sets and systems*, 2(4), 327-348.
- Imai, Y., & Iséki, K. (1965). On axiom systems of propositional calculi. I. *Proceedings of the Japan Academy*, 41(6), 436-439.
- Iséki, K. (1966). An algebra related with a propositional calculus. *Proceedings of the Japan Academy*, 42(1), 26-29.
- Jan, N., Maqsood, R., Nasir, A., Alhilal, M. S., Alabrah, A., & Al-Aidroos, N. (2022). A New Approach to Model Machine Learning by Using Complex Bipolar Intuitionistic Fuzzy Information. *Journal of Function Spaces*, 2022, 3147321.
- Jun, Y. B. (2005). On (α, β) -fuzzy subalgebras of BCK/BCI-algebras. *Bulletin of the Korean Mathematical Society*, 42(4), 703-711.
- Jun, Y. B. (2009). Generalizations of $(\epsilon, \epsilon \vee q)$ -fuzzy subalgebras in BCK/BCI-algebras. *Computers & Mathematics with Applications*, 58(7), 1383-1390.
- Jun, Y. B., & Xin, X. L. (2019). Complex fuzzy sets with application in BCK/BCI-Algebras. *Bulletin of the Section of Logic*, 48(3), 173-185.
- Jun, Y. B., Kang, M. S., & Kim, H. S. (2009). Bipolar fuzzy structures of some types of ideals in hyper BCK-algebras. *Scientiae Mathematicae Japonicae*, 70(1), 109-121.
- Kawila, K., Udomsetchai, C., & Iampan, A. (2018). Bipolar fuzzy UP-algebras. *Mathematical and Computational Applications*, 23(4), 69.
- Lee, K. J., & Jun, Y. B. (2011). Bipolar fuzzy a-ideals of BCI-algebras. *Communications of the Korean Mathematical Society*, 26(4), 531-542.
- Liu, W. J. (1982). Fuzzy invariant subgroups and fuzzy ideals. *Fuzzy sets and Systems*, 8(2), 133-139.
- Mahmood, T., & Munir, M. (2013). On bipolar fuzzy subgroups. *World Applied Sciences Journal*, 27(12), 1806-1811.
- Mahmood, T., & Ur Rehman, U. (2022a). A novel approach towards bipolar complex fuzzy sets and their applications in generalized similarity measures. *International Journal of Intelligent Systems*, 37(1), 535-567.
- Mahmood, T., & Ur Rehman, U. (2022b). A method to multi-attribute decision making technique based on Dombi aggregation operators under bipolar complex fuzzy information. *Computational and Applied Mathematics*, 41(1), 1-23.
- Mahmood, T., Rehman, U. U., Ahmmad, J., & Santos-García, G. (2021). Bipolar complex fuzzy Hamacher aggregation operators and their applications in multi-attribute decision making. *Mathematics*, 10(1), 23.
- Mahmood, T., Rehman, U. U., Jaleel, A., Ahmmad, J., & Chinram, R. (2022). Bipolar complex fuzzy soft sets and their applications in decision-making. *Mathematics*, 10(7), 1048.
- Meng, B. L. (2000). Some results of fuzzy BCK-filters. *Information Sciences*, 130(1-4), 185-194.
- Mordeson, J. N., & Malik, D. S. (1998). *Fuzzy commutative algebra*. Singapore, New Jersey, London, Hong Kong: World scientific.
- Muhiuddin, G. (2014). Bipolar fuzzy KU-subalgebras/ideals of KU-algebras. *Annals of Fuzzy Mathematics and Informatics*, 8(3), 409-418.
- Muhiuddin, G., & Al-Kadi, D. (2021). Bipolar fuzzy implicative ideals of BCK-algebras. *Journal of Mathematics*, 2021, 6623907.
- Muhiuddin, G., Al-Kadi, D., Mahboob, A., & Shum, K. P. (2020). New types of bipolar fuzzy ideals of BCK-algebras. *International Journal of Analysis and Applications*, 18(5), 859-875.
- Ramot, D., Milo, R., Friedman, M., & Kandel, A. (2002). Complex fuzzy sets. *IEEE Transactions on Fuzzy Systems*, 10(2), 171-186.
- Rasuli, R. (2022). Anti complex fuzzy Lie subalgebras under S-norms. *Mathematical Analysis and its Contemporary Applications*, 4(4), 13-26.
- Rehman, U. U., Mahmood, T., Albaity, M., Hayat, K., & Ali, Z. (2022). Identification and Prioritization of DevOps Success Factors Using Bipolar Complex Fuzzy Setting With Frank Aggregation Operators and Analytical Hierarchy Process. *IEEE Access*, 10, 74702-74721.

- Rezaei, A., & Saeid, A. B. (2011). On fuzzy subalgebras of BE-algebras. *Afrika Matematika*, 22(2), 115-127.
- Shaqqa, S. (2019). Complex fuzzy lie algebras. arXiv preprint arXiv:1905.01687.
- Tamir, D. E., Jin, L., & Kandel, A. (2011). A new interpretation of complex membership grade. *International Journal of Intelligent Systems*, 26(4), 285-312.
- Yang, X., Mahmood, T., & ur Rehman, U. (2022). Bipolar Complex Fuzzy Subgroups. *Mathematics*, 10(16), 2882.
- Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8(3), 338-353.
- Zhang, W. R. (1994). Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis. In *NAFIPS/IFIS/NASA'94. Proceedings of the First International Joint Conference of The North American Fuzzy Information Processing Society Biannual Conference. The Industrial Fuzzy Control and Intellige, IEEE*, 305-309.